




2007 Carbon Monoxide Summary

New Jersey Department of Environmental Protection

NATURE AND SOURCES

Carbon monoxide (CO) is a colorless, odorless, poisonous gas formed when carbon in fuels is not burned completely. It is a by-product of motor vehicle exhaust, which contributes over 56 percent of all CO emissions nationwide. In cities, automobile exhaust can cause as much as 95 percent of all CO emissions, and high CO levels often coincide with morning and afternoon rush hours (Figure 4 on page 3). Non-road engines and vehicles, such as construction equipment and boats, are also significant sources of CO and overall the transportation sector is responsible for about 82% of all CO emissions nationally. Other sources of CO include industrial processes, fuel combustion in sources such as boilers and incinerators, and natural sources such as forest fires. Figure 1 shows the national average contributions of these sources.

**Figure 1
National Summary of
CO Emissions by Source Category**





Source: United States Environmental Protection Agency
www.epa.gov/air/urbanair/co/what1.html
Updated: July, 2007

Figure 3 shows that CO levels are slightly higher in the winter. This is because motor vehicles do not burn fuel as efficiently when they are cold. Atmospheric inversions are also more frequent during the winter months. Inversions usually occur overnight when cooler air is trapped beneath a layer of warmer air aloft. When this occurs, the inversion acts

like a lid, preventing pollution from mixing in the atmosphere and effectively trapping it close to ground level (see Figure 2).

Figure 2: Effect of Atmospheric Inversion on Air Pollution



HEALTH AND ENVIRONMENTAL EFFECTS

Carbon monoxide enters the bloodstream and reduces the body's ability to distribute oxygen to organs and tissues. The most common symptoms associated with exposure to carbon monoxide are headaches and nausea. The health threat from exposure to CO is most serious for those who suffer from cardiovascular disease. For a person with

heart disease, a single exposure to CO at low levels may cause chest pain and reduce that individual's ability to exercise. Healthy people are also affected, but only at higher levels of exposure. Elevated CO levels are also associated with visual impairment, reduced work capacity, reduced manual dexterity, decreased learning ability, and difficulty in performing complex tasks.

opposed to parts per million), and our standards are not to be exceeded more than once in any 12-month period. The state has set secondary (welfare based) standards for CO at the same level as the primary standards. The standards are summarized in Table 1.

STANDARDS

There are currently two national primary, or health based, standards for carbon monoxide. They are set at a one-hour concentration of 35 parts per million (ppm), and an 8-hour average concentration of 9 ppm. These levels are not to be exceeded more than once in any calendar year. There are no national secondary (welfare based) standards for CO at this time.

New Jersey state standards for CO are based on different units (milligrams per cubic meter as

Table 1
National and New Jersey Ambient Air Quality Standards for Carbon Monoxide

mg/m³ = Milligrams Per Cubic Meter
ppm = Parts per Million

Averaging Period	Type	New Jersey	National
1-Hour	Primary	40 mg/m ³ (35 ppm)	35 ppm
1-Hour	Secondary	40 mg/m ³ (35 ppm)	----
8-Hour	Primary	10 mg/m ³ (9 ppm)	9 ppm
8-Hour	Secondary	10 mg/m ³ (9 ppm)	----

Figure 3
2007 Carbon Monoxide Concentrations – New Jersey
Monthly Variation
Parts Per Million (PPM)





Figure 4
2007 Carbon Monoxide Concentrations – New Jersey
Diurnal Variation
Parts Per Million (PPM)



MONITORING LOCATIONS

The state monitored CO levels at 12 locations in 2007. These sites are shown in the map in Figure 5. The Elizabeth site was temporarily shut down starting in March 2007 through the end of the year.

CO LEVELS IN 2007

None of the monitoring sites recorded exceedances of any CO standard during 2007. The maximum one-hour average concentration recorded was 3.4 ppm at the site in Elizabeth Lab. The highest 8-hour average level recorded was 2.7 ppm, recorded at the East Orange site. Summaries of the 2007 data are provided in Figure 6 and Table 2 (page 4).

Figure 5
2007 Carbon Monoxide
Monitoring Network



Figure 6
Highest and 2nd Highest 8-Hour Averages
Of Carbon Monoxide in New Jersey – 2007
Parts Per Million (PPM)




Table 2
Carbon Monoxide Data – 2007
1-Hour and 8-Hour Averages

Parts Per Million (ppm)
 1-hour standard = 35 ppm
 8-hour standard = 9 ppm

Monitoring Sites	Maximum 1-Hour Average	2 nd Highest 1-Hour Average	Maximum 8-Hour Average	2 nd Highest 8-Hour Average
Ancora State Hospital	1.7	1.3	0.7	0.6
Burlington	2.4	2.3	1.4	1.4
Camden Lab	2.9	2.5	2.1	2.0
East Orange	3.2	3.2	2.7	2.1
Elizabeth*	2.9	2.7	1.9	1.8
Elizabeth Lab	3.4	2.9	2.6	1.9
Fort Lee	2.3	2.2	1.8	1.5
Freehold	3.3	2.0	1.7	1.4
Hackensack	2.3	2.2	1.8	1.8
Jersey City	3.3	2.6	2.1	1.9
Morristown	2.2	2.1	1.2	1.2
Perth Amboy	2.4	2.0	1.6	1.4

* Temporary shut down of site starting March 2007

Trends

Carbon monoxide levels have improved dramatically over the past 20 years. The last time the CO standard was exceeded in New Jersey was in January of 1995 (Figure 7), and the entire state was officially declared as having attained the CO standard on August 23, 2002. At one time, unhealthy levels of CO were recorded on a regular basis – more than a hundred days a year at some sites. The reduction in CO levels is due primarily to cleaner running cars, which are by far the largest source of this pollutant. A trend graph of CO levels showing the concentrations of the Highest Site, Average of All Sites, and Lowest Site of each year since 1975 is provided in Figure 8. The graph depicts the second highest 8-hour value recorded as this is the value that determines if the health standard is being met (one exceedance per site is allowed each year).

Figure 7
Carbon Monoxide
Unhealthy Days 1985-2007





Figure 8
Carbon Monoxide Air Quality, 1975-2007
2nd Highest 8-Hour Average
Parts Per Million (PPM)



REFERENCES

CO – How Carbon Monoxide Affects the Way We Live and Breathe, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC November 2000, URL: <http://www.epa.gov/air/urbanair/co/index.html>

Automobile Emissions: An Overview, USEPA, August 1994, EPA-400/F-92-007, URL: <http://www.epa.gov/otaq/consumer/05-autos.pdf>

National Air Quality and Emissions Trend Report, 1999, EPA-454/R-01-004, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, March 2001, URL: <http://www.epa.gov/air/airtrends/aqtrnd99/>

Latest Findings on National Air Quality: 2000 Status and Trends, EPA-454/K-01-002, USEPA, Office of Air Quality Planning and Standards, RTP, September 2001, URL: <http://www.epa.gov/air/airtrends/aqtrnd00/brochure/00brochure.pdf>

Latest Findings on National Air Quality: 2002 Status and Trends, EPA-454/K-03-001, USEPA, Office of Air Quality Planning and Standards, RTP, September 2001, URL: www.epa.gov/airtrends/carbon.html

National Air Quality and Emissions Trend Report, 2003 Special Studies Edition, EPA-454/R-03-005, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, September 2003, URL: <http://www.epa.gov/air/airtrends/aqtrnd03/>

National Primary Ambient Air Quality Standards for Carbon Monoxide, 40 CFR 50.8, US Government Printing Office, Washington DC, July 2001.