




# 2010 Nitrogen Dioxide Summary

New Jersey Department of Environmental Protection

## NATURE AND SOURCES

Nitrogen Dioxide ( $\text{NO}_2$ ) is a reddish-brown, highly reactive gas that is formed in the air through the oxidation of Nitric Oxide (NO). When  $\text{NO}_2$  reacts with other chemicals, it can form ozone, particulate matter, and other compounds which can contribute to regional haze and acid rain. Oxides of Nitrogen ( $\text{NO}_x$ ) is a mixture of gases which is mostly comprised of NO and  $\text{NO}_2$ . These gases are emitted from the exhaust of motor vehicles, the burning of coal, oil or natural gas, and during industrial processes such as welding, electroplating, and dynamite blasting. Although most  $\text{NO}_x$  is emitted as NO, it is readily converted to  $\text{NO}_2$  in the atmosphere. In the home, gas stoves and heaters produce substantial amounts of nitrogen dioxide. A pie chart summarizing the major sources of  $\text{NO}_x$  is shown below (Figure 1). As much of the  $\text{NO}_x$  in the air is emitted by motor vehicles, concentrations tend to peak during the morning and afternoon rush hours. This is shown in the graph in Figures 2-4 (pages 2-3). Figures 6-8 (pages 6-7) indicate that concentrations tend to be higher in the winter than the summer. This is due in part to space heating and poorer local dispersion conditions caused by light winds and other weather conditions that are more prevalent in the colder months of the year.

**Figure 1**  
**National Summary of 2005 Oxides of Nitrogen ( $\text{NO}_x$ ) Emissions by Source Category**



Source: USEPA National Summary of Nitrogen Oxides Emissions, 2005

## HEALTH AND ENVIRONMENTAL EFFECTS

Short-term exposures (less than 3 hours) to low levels of nitrogen dioxide may aggravate pre-existing respiratory illnesses, and can cause respiratory illnesses, particularly in children ages 5-12. Symptoms of low level exposure to NO and  $\text{NO}_2$  include irritation to eyes, nose, throat and lungs, coughing, shortness of breath, tiredness and nausea. Long-term exposures to  $\text{NO}_2$  may increase susceptibility to respiratory infection and may cause permanent damage to the lung. NO and  $\text{NO}_2$  are found in tobacco smoke, so people who smoke or breathe in second-hand smoke may be exposed to  $\text{NO}_x$ . The U.S. Department of Health and Human Services (DHHS), the International Agency for Research on Cancer (IARC), and the U.S. Environmental Protection Agency (EPA) have determined that, with the available information, no conclusion can be made as to the carcinogenicity of NO or  $\text{NO}_2$  to human beings.

Nitrogen Oxides contribute to a wide range of environmental problems. These include potential changes in the composition of some plants in wetland and terrestrial ecosystems, acidification of freshwater bodies, eutrophication of estuarine and coastal waters, increases in levels of toxins harmful to fish and other aquatic life, and visibility impairment.


## STANDARDS

The primary (health based) and secondary (welfare based) National Ambient Air Quality Standards (NAAQS) annual average for  $\text{NO}_2$  are the same. They are set at a calendar year average concentration of 0.053 parts per million (ppm). The New Jersey Ambient Air Quality Standards (NJAAQS) are identical to the NAAQS except micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ) are the standard units and the state standard applies to any 12-month period, not just the calendar year. In 2010, the EPA strengthened the primary NAAQS by adding a 1-hour  $\text{NO}_2$  standard of 0.100 ppm along with the current annual average  $\text{NO}_2$  standard of 0.053 ppm. Table 1 provides a summary of the  $\text{NO}_2$  standards.


**Table 1**  
**National and New Jersey Ambient Air Quality Standards for Nitrogen Dioxide (NO<sub>2</sub>)**  
**Parts Per Million (ppm) and Micrograms Per Cubic Meter (µg/m<sup>3</sup>)**

| Averaging Period | Type      | New Jersey                        | National                           |
|------------------|-----------|-----------------------------------|------------------------------------|
| 12-month average | Primary   | 100 µg/m <sup>3</sup> (0.053 ppm) |                                    |
| Annual average   | Primary   |                                   | 0.053 ppm (100 µg/m <sup>3</sup> ) |
| 12-month average | Secondary | 100 µg/m <sup>3</sup> (0.053 ppm) |                                    |
| Annual average   | Secondary |                                   | 0.053 ppm (100 µg/m <sup>3</sup> ) |
| 1-hour average   | Primary   |                                   | 0.100 ppm (190 µg/m <sup>3</sup> ) |


**Figure 2**  
**Nitric Oxide – New Jersey**  
**2010 Hourly Variation**  
**Parts Per Million (ppm)**



**Figure 3**  
**Nitrogen Dioxide – New Jersey**  
**2010 Hourly Variation**  
**Parts Per Million (ppm)**



**Figure 4**  
**Total Oxides of Nitrogen – New Jersey**  
**2010 Hourly Variation**  
**Parts Per Million (ppm)**



## STANDARDS (CONTINUED)

An area meets the new 1-hour NO<sub>2</sub> standard when the 3-year average of the 98<sup>th</sup> percentile of the daily maximum NO<sub>2</sub> concentrations measured in this area is less than 0.100 ppm. This statistic, also known as the design value, is determined by first obtaining the maximum 1-hour average NO<sub>2</sub> concentrations for each day. Then, determine the 98<sup>th</sup> percentile of the daily maximum NO<sub>2</sub> concentrations for the current year, and for each of the previous two years. Finally, the average of these three 98<sup>th</sup> percentile values is the design value.

## MONITORING LOCATIONS

The state monitored NO<sub>2</sub> levels at 8 locations in 2010. These sites are shown in the map on page 5. The NO<sub>2</sub> analyzers at the Leonia and Rider University sites were shut down on December 31, 2010.


## NO<sub>2</sub> LEVELS IN 2010

None of the monitoring sites recorded exceedances of either the National or New Jersey Air Quality Standards for NO<sub>2</sub> during 2010. The highest 12-month (calendar year) average concentration of NO<sub>2</sub> recorded was 0.022 ppm at the Elizabeth Lab site located at Exit 13 of the New Jersey Turnpike (Table 2, page 6 and Figure 9, page 8). At Bayonne, there was one measurement of the 1-hour average NO<sub>2</sub> concentration above 0.100 ppm. This occurred at Elizabeth Lab on a different day. The site that measured the highest 98<sup>th</sup> percentile of the daily maximum NO<sub>2</sub> concentrations in 2010 was Elizabeth Lab with 0.071 ppm (Table 2, page 6 and Figure 13, page 10). The site that measured the highest 3-year average of the 98<sup>th</sup> percentiles from 2008 to 2010 was also Elizabeth Lab with 0.073 ppm (Table 2, page 6 and Figure 12, page 9). All sites in New Jersey met the new 1-hour NO<sub>2</sub> standard. While national health and welfare standards have not been established for Nitric Oxide (NO), it is considered to be an important pollutant that contributes to the formation of ozone, fine particles and acid rain. The maximum annual average concentration of NO recorded in 2010 was 0.021 ppm, also at the Elizabeth Lab site (Table 2, page 6 and Figure 10, page 8).

## TRENDS

Routine monitoring for NO<sub>2</sub> began in 1966 and 1974 was the last year that concentrations exceeded the NAAQS in New Jersey. A graph of NO<sub>2</sub> levels provided in Figure 11 (page 9) shows the statewide average annual mean concentrations recorded from 1975 to 2010 in the form of a trendline. The graph also includes the levels of the sites that measured the highest annual mean and lowest annual mean in each year as points above and below this trendline. Although NO<sub>2</sub> concentrations are well within the NAAQS, there is still a great deal of interest in oxides of nitrogen because of their role in the formation of other pollutants – most notably ozone and fine particles. Both these pollutants are of concern over much of the northeastern United States and efforts to reduce levels of ozone and fine particles are likely to require reductions in NO emissions.

**Figure 5**  
**2010 Nitrogen Dioxide**  
**Monitoring Network**




**Table 2**  
**Nitrogen Dioxide (NO<sub>2</sub>) and Nitric Oxide (NO) Data - 2010**  
**1-Hour and 12-Month Averages**


Parts Per Million (ppm)  
 National 1-Hour Standard = 0.100 ppm  
 National 12-Month Standard = 0.053 ppm

| Monitoring Sites   | Nitrogen Dioxide     |             |                   |   | Nitrogen Dioxide                  |               | Nitric Oxide  |
|--------------------|----------------------|-------------|-------------------|---|-----------------------------------|---------------|---------------|
|                    | 1-Hour Average (ppm) |             |                   |   | 12-Month Average (ppm)            |               | 12-Month      |
|                    | Maximum              | 2nd Highest | 2010<br>98th%-ile | 2008-2010<br>98 <sup>th</sup> %-ile<br>(3-year) | Maximum<br>(Running 12-<br>month) | Calendar year | Average (ppm) |
|                    |                      |             |                   |   |                                   |               | Calendar Year |
| Bayonne            | 0.106                | 0.096       | 0.068             | 0.067   | 0.018                             | 0.018         | 0.009         |
| Chester            | 0.063                | 0.049       | 0.035             | 0.038   | 0.004                             | 0.004         | 0.000         |
| East Orange        | 0.072                | 0.072       | 0.064             | 0.062   | 0.019                             | 0.018         | 0.012         |
| Elizabeth Lab      | 0.108                | 0.094       | 0.071             | 0.073   | 0.026                             | 0.022         | 0.021         |
| Leonia             | 0.080                | 0.071       | 0.064             | 0.066   | 0.017                             | 0.015         | 0.013         |
| Millville          | 0.054                | 0.050       | 0.040             | 0.040   | 0.009                             | 0.008         | 0.005         |
| Rider University   | 0.046                | 0.045       | 0.040             | 0.041   | 0.008                             | 0.008         | 0.004         |
| Rutgers University | 0.052                | 0.049       | 0.045             | 0.049   | 0.011                             | 0.010         | 0.003         |


**Figure 6**  
**Nitric Oxide – New Jersey**  
**2010 Monthly Variation**  
**Parts Per Million (ppm)**




**Figure 7**  
**Nitrogen Dioxide – New Jersey**  
**2010 Monthly Variation**  
**Parts Per Million (ppm)**




**Figure 8**  
**Total Oxides of Nitrogen – New Jersey**  
**2010 Monthly Variation**  
**Parts Per Million (ppm)**



**Figure 9**  
**Annual Average Nitrogen Dioxide Concentrations**  
**In New Jersey – 2010**  
**Parts Per Million (ppm)**




**Figure 10**  
**Annual Average Nitric Oxide Concentrations**  
**In New Jersey – 2010**  
**Parts Per Million (ppm)**







**Figure 11**  
**Nitrogen Dioxide Concentrations in New Jersey 1975-2010**  
**12-Month (Calendar Year) Average**  
**Parts Per Million (ppm)**



**Figure 12**  
**3-Year Average 98<sup>th</sup> Percentile Nitrogen Dioxide Concentration**  
**In New Jersey (2008-2010)**  
**Parts Per Million (ppm)**



**Figure 13**  
**2010 Average 98<sup>th</sup> Percentile Nitrogen Dioxide Concentration**  
**In New Jersey**  
**Parts Per Million (ppm)**



## REFERENCES

*Fact Sheet Final Revisions to the National Ambient Air Quality Standards for Nitrogen Dioxide*, USEPA, January 22, 2010, URL: <http://www.epa.gov/airquality/nitrogenoxides/pdfs/20100122fs.pdf>.

*History of California's Ambient Air Quality Standards*, California Air Resources Board (CARB), Sacramento, California, March 5, 2008, URL: <http://www.arb.ca.gov/research/aaqs/caaqs/hist1/hist1.htm>

Meyer, Edwin L., Sennet, Donald H., Cole, Henry S., Richter, Harold G., *Technical Basis for Developing Control Strategies for High Ambient Concentrations of Nitrogen Dioxide*, EPA-450/4-80-017, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, 1980.

*National Primary and Secondary Ambient Air Quality Standards for Nitrogen Dioxide*, 40 CFR 50.11, US Government Printing Office, Washington DC, July 2001.

*National Summary of Nitrogen Oxides Emissions*, USEPA, Research Triangle Park, NC, 2005, URL: <http://www.epa.gov/air/emissions/nox.htm>.

*Nitrogen Dioxide and Respiratory Illness in Children*, Health Effects Institute, 1994.

*NO<sub>x</sub> – How Nitrogen Oxides Affect the Way We Live and Breathe*, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, September 1998, URL: <http://www.epa.gov/air/nitrogenoxides/>.

*The Regional Transport of Ozone, New EPA Rulemaking on Nitrogen Oxide Emissions*, EPA-456/F-98-006, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, URL: <http://www.epa.gov/air/noxfacts.pdf>.

*Review Of The National Ambient Air Quality Standards For Nitrogen Dioxide Assessment Of Scientific And Technical Information*, EPA-452/R-95-005, OAQPS staff paper, USEPA, Office of Air and Radiation, Office of Air Quality Planning and Standards, 1995.

Sittig, M., *Handbook of Toxic and Hazardous Chemicals and Carcinogens Third Edition*, Volume 2, Noyes Publications, Park Ridge, NJ, 1991.

*ToxFaQs for Nitrogen Oxides*, U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, April 2002, URL: <http://www.atsdr.cdc.gov/tfacts175.pdf>.

Utell, Mark J., *Mechanisms of Nitrogen Dioxide Toxicity in Humans*, Health Effects Institute, 1991