



2013 Ozone Summary

New Jersey Department of Environmental Protection


NATURE AND SOURCES

Ozone (O₃) is a gas consisting of three oxygen atoms. It occurs naturally in the upper atmosphere (stratospheric ozone) where it protects us from harmful ultraviolet rays (see Figure 1). However, at ground-level (tropospheric ozone) it is considered an air pollutant and can have serious adverse health effects. Ground-level ozone is created when nitrogen oxides (NO_x) and volatile organic compounds (VOC's) react in the presence of sunlight. NO_x is primarily emitted by motor vehicles, power plants, and other sources of combustion. VOC's are emitted from sources such as motor vehicles, chemical plants, factories, consumer and commercial products, and even natural sources such as trees. Ozone and the pollutants that form ozone (precursor pollutants) can also be transported into an area from sources hundreds of miles upwind.

Since ground-level ozone needs sunlight to form, it is mainly a daytime problem during the summer months. Weather patterns have a significant effect on ozone formation and hot, dry summers will result in more ozone than cool, wet ones. In New Jersey, the ozone monitoring season runs from April 1st to October 31st. For a more complete explanation of the difference between ozone in the upper and lower atmosphere, see the U.S. Environmental Protection Agency (EPA) publication "Ozone: Good Up High, Bad Nearby".

Figure 1: Good and Bad Ozone

OZONE IS GOOD UP HERE...MANY POPULAR CONSUMER PRODUCTS LIKE AIR CONDITIONERS AND REFRIGERATORS INVOLVE CFCs OR HALONS DURING EITHER MANUFACTURING OR USE. OVER TIME, THESE CHEMICALS DAMAGE THE EARTH'S PROTECTIVE OZONE LAYER.




OZONE IS BAD DOWN HERE... CARS, TRUCKS, POWER PLANTS AND FACTORIES ALL EMIT AIR POLLUTION THAT FORMS GROUND-LEVEL OZONE, A PRIMARY COMPONENT OF SMOG.

ENVIRONMENTAL EFFECTS

Ground-level ozone damages plant life and is responsible for 500 million dollars in reduced crop production in the United States each year. It interferes with the ability of plants to produce and store food, making them more susceptible to disease, insects, other pollutants, and harsh weather. "Bad" ozone damages the foliage of trees and other plants, sometimes marring the landscape of cities, national parks and forests, and recreation areas. The black areas on the leaves of the watermelon plant, shown in Figure 2, are damage caused by exposure to ground-level ozone.

Figure 2 - Damage Caused by Ozone




(Photos by: Gerald Holmes, NCSU Dept. of Horticulture)

HEALTH EFFECTS

Repeated exposure to ozone pollution may cause permanent damage to the lungs. Even when ozone is present in low levels, inhaling it can trigger a variety of health problems including chest pains, coughing, nausea, throat irritation, and congestion. Ozone also can aggravate other health problems such as bronchitis, heart disease, emphysema, and asthma, and can reduce lung capacity. People with pre-existing respiratory ailments are especially prone to the effects of ozone. For example, asthmatics affected by ozone may have more frequent or severe attacks during periods when ozone levels are high. As shown in Figure 3 ozone can irritate the entire respiratory tract. Children are also at risk for ozone related problems. Their respiratory systems are still developing and they breathe more air per pound of body weight than adults. They are also generally active outdoors during the summer when ozone levels are at their highest. Anyone who spends time outdoors in the summer can be affected and studies have shown that even healthy adults can experience difficulty in breathing when exposed to ozone. Anyone engaged in strenuous outdoor activities, such as jogging, should limit activity to the early morning or late evening hours on days when ozone levels are expected to be high.

Figure 3
Effects of Ozone & Common Air Pollutants



Source: www.airnow.gov

AMBIENT AIR QUALITY STANDARDS FOR OZONE

National and state air quality standards have been established for ground-level ozone. There are both primary standards, which are based on health effects, and secondary standards, which are based on welfare effects (e.g. damage to trees, crops and materials). For ground-level ozone, the primary and secondary National Ambient Air Quality Standards (NAAQS) are the same (see Table 1). The ozone NAAQS were revised in 2008 because EPA determined that the old standard of 0.08 parts per million (ppm) maximum daily eight-hour average was not sufficiently protective of public health. The revised standard of 0.075 ppm maximum daily 8-hour average went into effect on May 27, 2008.

As many people are accustomed to the old standards, summary information relative to that standard will be provided in this report along with summaries based on the new standard.

Table 1
National and New Jersey Ambient Air Quality Standards for Ozone


ppm = Parts per Million

Averaging Period	Type	New Jersey	National
1-Hour	Primary	0.12 ppm	-----
1-Hour	Secondary	0.08 ppm	-----
8-Hour	Primary	-----	0.075 ppm
8-Hour	Secondary	-----	0.075 ppm

OZONE NETWORK

Ozone was monitored at 16 locations in New Jersey during 2013. (See Figure 4) Of those 16 sites, 10 operated year round and 6 operated only during the ozone season (April 1st through October 31st). Ancora State Hospital, Clarksboro, Colliers Mills, Leonia, Monmouth University, and Ramapo were only operated during the ozone season.

Figure 4
2013 Ozone Monitoring Network




HOW THE CHANGES TO THE OZONE STANDARDS AFFECT AIR QUALITY RATINGS

In May of 2008 the U.S. Environmental Protection Agency revised the NAAQS for ozone from a daily maximum 8-hour average concentration of 0.08 ppm to a daily maximum 8-hour average of 0.075 ppm. While this may not seem like a major change, it does result in significantly more days with levels above the standard recorded. In 2013 for example, there were 10 days on which the 0.075 ppm was exceeded, but only 2 days on which the old 0.08 ppm standard was exceeded.

However, exceedances of both standards are still recorded on a regular basis (see Figure 5 below). As a result, additional control measures to reduce ozone levels will be needed. These measures will have to be implemented over a wide area and will require the cooperative effort of many states and the federal government if they are to be successful. In figure 5 the new standard has been projected back through 2000 for comparison purposes.

Figure 5
Days on Which the Old and New
Ozone Standards have been exceeded in New Jersey
1993-2013



DESIGN VALUE

The NAAQS for ozone are set in such a way that determining whether they are being attained is not based on a single year. For example, an area was considered to be attaining the old 1-hour average standard if the average number of times the standard was exceeded over a three-year period was 1 or less (after correcting for missing data). Thus it was the fourth highest daily maximum 1-hour concentration that occurred over a three-year period that determined if an area would be in attainment. If the fourth highest value was above 0.12 ppm then the average number of exceedances would be greater than 1. The fourth highest value is also known as the design value.

Under the new standard, attainment is determined by taking the average of the fourth highest daily maximum 8-hour average concentration that is recorded each year for three years. This becomes the design value for an area under the new standard. When plans are developed for reducing ozone concentrations, an area must demonstrate that the ozone reduction achieved will be sufficient to ensure the design value will be below the NAAQS, as opposed to ensuring that the standards are never exceeded. This avoids developing plans based on extremely rare events.

Table 2 and Table 3 on the following pages display the current design values for the 1-hour standard and the 8-hour standard respectively.

SUMMARY OF 2012 OZONE DATA RELATIVE TO THE OLD 1-HOUR STANDARD

Of the 16 monitoring sites that were operated during the 2013 ozone season, none recorded levels above the old 1-hour standard of 0.12 ppm. The highest 1-hour concentration was 0.105 ppm recorded at Bayonne and Monmouth University on July 17th and 18th, respectively. As recently as 2002, New Jersey recorded 16 days above this old 1-hour standard. Figure 6 on the following page shows both the highest and second highest daily 1-hour averages.


Table 2
Ozone Data – 2013
1-Hour Averages
 Parts Per Million (ppm)

Old 1-hour standard is 0.12 ppm

Monitoring Site	1-hr Max	2nd Highest 1-hr Max	4th Highest 1-hour Average 2011-2013	# of days in 2013 with 1-hour Average above 0.12ppm
Ancora S.H.	0.083	0.081	0.105	0
Bayonne	0.105	0.100	0.102	0
Brigantine	0.103	0.096	0.096	0
Camden Spruce St.*	0.084	0.084	---	0
Chester	0.092	0.089	0.093	0
Clarksboro	0.085	0.084	0.107	0
Colliers Mills	0.089	0.088	0.107	0
Columbia WMA	0.094	0.089	0.092	0
Flemington	0.095	0.094	0.095	0
Leonia	0.099	0.093	0.101	0
Millville	0.078	0.077	0.097	0
Monmouth Univ.	0.105	0.101	0.105	0
Newark Firehouse	0.084	0.083	0.100	0
Ramapo	0.093	0.088	0.091	0
Rider University	0.085	0.083	0.101	0
Rutgers University	0.089	0.085	0.104	0

*Camden Spruce St. data only available for 2012-2013.

Figure 6
 New Jersey Ozone Data - 2013
 Highest and Second Highest
 Daily Maximum 1-hour Averages



Summary of 2013 Ozone Data Relative to the 8-Hour Standard

Just 11 of 16 monitoring sites operating during the 2013 ozone season recorded levels above the 8-hour standard of 0.075 ppm. The highest daily maximum 8-hour concentration recorded was 0.088 ppm at Brigantine on July 17th. Design values for the 8-hour standard were above the standard at 11 of 16 sites, indicating that the ozone standard is being violated throughout almost all of New Jersey. Camden Spruce St. does not have three complete years of data to calculate a valid design value. Figure 7 on the following page charts the 8-hour design values for the 2011-2013 period.

Table 3
Ozone Data – 2013
8-Hour Averages
 Parts Per Million (ppm)

Monitoring Site	Highest Daily Maximum 8-hour Average for 2013	Average of 4 th Highest Daily Maximum 8-hour Averages, 2011-2013	# of days in 2013 with 8-hour Average above 0.075 ppm
Ancora S.H.	.075	0.081	0
Bayonne	.086	0.072	1
Brigantine	.088	0.073	2
Camden Spruce St.*	.073	---	---
Chester	.078	0.076	2
Clarksboro	.079	0.084	2
Colliers Mills	.076	0.080	1
Columbia WMA	.076	0.066	1
Flemington	.077	0.077	1
Leonia	.077	0.077	2
Millville	.068	0.070	0
Monmouth Univ.	.087	0.078	3
Newark Firehouse	.079	0.076	2
Ramapo	.078	0.072	2
Rider University	.075	0.076	0
Rutgers University	.075	0.079	0

**Camden Spruce St. only has 2 years of data and does not have a valid design value for 2011-2013.*

Figure 7
 Ozone Design Values for 2011-2013
 3-Year Average of the 4th Highest Daily Maximum 8-hour Average





Figure 8
 Number of Days 8-Hour Ozone Standard was Exceeded and
 Number of Days Above 90 Degrees in New Jersey 1988-2013
 (Using 8-Hour 0.08 ppm standard (old) across entire time period)




ACCOUNTING FOR THE INFLUENCE OF WEATHER

Trends in ground level ozone are influenced by many factors including weather conditions, transport, growth, and the state of the economy, in addition to changes brought about by regulatory control measures. Of these factors, weather probably has the most profound effect on year to year variations in ozone levels. Several methods have been developed to try to account for the effect of weather on ozone levels so that the change due to emissions could be isolated. While none of these methods are completely successful they do show that over the long term, real reductions in ozone levels have been achieved. A simple way of showing the changing effect of weather on ozone is shown above in Figure 8. The number of days each year on which the ambient temperature was 90 degrees or greater is shown next to the number of days the ozone standard was exceeded. In the earliest years shown (1988-1993) there are significantly more days with high ozone than days above 90 degrees. But this pattern gradually changes and for the most recent years there are more “hot” days than “ozone” days. This is an indication that on the days when conditions are suitable for ozone formation, unhealthy levels are being reached less frequently.

OZONE TRENDS

The primary focus of efforts to reduce concentrations of ground-level ozone in New Jersey has been on reducing emissions of volatile organic compounds (VOCs). Studies have shown that such an approach should lower peak ozone concentrations, and it does appear to have been effective in achieving that goal. The chart in Figure 9 is based on the second highest 1-hour average concentrations recorded each year. We use this statistic when showing long term trends as it is what the early ozone health standards were based on, so historical data including those values is readily available. As Figure 9 illustrates, the maximum 1-hour concentrations have not exceeded 0.200 ppm since 1988 and the last time levels above 0.180 ppm were recorded was in 1990. Improvements have leveled off in recent years; and further improvements will require reductions in both VOCs and NO_x. The NO_x reductions will have to be achieved over a very large region of the country because levels in New Jersey are dependent on emissions from upwind sources.



OZONE NON-ATTAINMENT AREAS IN NEW JERSEY


The Clean Air Act requires that all areas of the country be evaluated and then classified as attainment or non-attainment areas for each of the National Ambient Air Quality Standards. Areas can also be found to be “unclassifiable” under certain circumstances. The 1990 amendments to the act required that areas be further classified based on the severity of non-attainment. The classifications range from “Marginal” to “Extreme” and are based on “design values”. The design value is the value that actually determines whether an area meets the standard. For the 8-hour ozone standard for example, the design value is the average of the fourth highest daily maximum 8-hour average concentration recorded each year for three years.

Their classification with respect to the 8-hour standard is shown in Figure 10 below. The entire state of New Jersey is in non-attainment and is classified as being “Marginal.” A “Marginal” classification is applied when an area has a design value of 0.085 ppm up to but not including 0.092 ppm.

Figure 10

New Jersey 8-hour Ozone Nonattainment Areas (2008 Standard)

7/02/2014



Source: U.S. EPA Greenbook

REFERENCES

Ozone: Good Up High, Bad Nearby, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC October 1997, URL: www.epa.gov/oar/oaqps/gooduphigh/

USEPA Fact Sheet: Health and Environmental Effects of Ground Level Ozone, USEPA, Office of Air and Radiation, July 1997, URL: www.epa.gov/ttn/oarpg/naaqsf/o3health.html

Guidelines for Developing an Air Quality (Ozone and PM2.5) Forecasting Program, EPA-456/R-03-002, June 2003

National Air Quality and Emissions Trend Report, 1999, EPA-454/R-01-004, USEPA, Office of Air Quality Planning and Standards, Research Triangle Park, NC, March 2001, URL: www.epa.gov/airtrends/aqtrnd99/

Latest Findings on National Air Quality: 2000 Status and Trends, EPA-454/K-01-002, USEPA, Office of Air Quality Planning and Standards, RTP, September 2001, URL: www.epa.gov/airtrends/aqtrnd00/brochure/00brochure.pdf

Smog – Who Does it Hurt?, EPA-452/K-99-001, USEPA, Air and Radiation, Washington, DC, July 1999, URL: www.epa.gov/airnow/health/smog.pdf

Ozone and Your Health, EPA-152/F-99-000, USEPA, Air and Radiation, Washington, DC, September 1999, URL: www.epa.gov/airnow/ozone-c.pdf

Air Quality Guide for Ozone, EPA-456/F-002, Air and Radiation, Washington, DC, July 1999, URL: www.airnow.gov/index.cfm?action=aqibasics.aqi

Archived temperature data gathered from NOAA/NWS URL: www.weather.gov/climate/index.php?wfo=phi

USEPA Greenbook, URL: www.epa.gov/oar/oaqps/greenbk/nj8_2008.html